

IMPROVEMENT OF ENGINEERED THERMOSTABLE XYLANASE
PRODUCTION VIA IMMOBILIZATION OF RECOMBINANT
ESCHERICHIA COLI ONTO GRAPHENE OXIDE

NUR ATIQA LYANA BINTI NOR ASHIKIN

A thesis submitted in
fulfillment of the requirement for the award of the
Degree of Master of Science

Faculty of Applied Sciences and Technology
Universiti Tun Hussein Onn Malaysia

AUGUST 2020

For my beloved husband (Nuur Muhaimin Bin Hussin), parents (Nor Ashikin Bin Abu Bakar and Siti Junaidah Bt Ithnin), my brothers (Mohd Syafiq Faiz Bin Nor Ashikin, Muhammad Syarafi Iqbal Bin Nor Ashikin and Muhammad Safiy Haziq Bin Nor Ashikin), Sister (Nurin Syahindah Syasya Binti Nor Ashikin), I dedicated this work in sincere gratitude for their patience, love and support.



ACKNOWLEDGEMENT

The author would like to express her sincere appreciation to her supervisor, Dr. Siti Fatimah Zaharah Binti Md Fuzi, co-supervisor; Prof. Dr. Rosli bin Md Illias, Dr Low Kheng Oon for the support given throughout the duration for this research.

The high appreciation goes to the Ministry of Higher Education (MOHE) and Universiti Tun Hussien Onn Malaysia (UTHM) for providing the scholarship and allowance during the study.

The cooperation given by the Malaysia Genome Institute staffs genetic laboratory members at Universiti Teknologi Malaysia (UTM) are also highly appreciated.

Appreciation also goes to everyone involved directly or indirectly towards the compilation of this thesis. Last but not least, special appreciation to her family and friends for the patience, understanding and support through ups and downs, good and bad in order to help in accomplishment of this project.

Thank you very much.



ABSTRACT

Escherichia coli is the preference host system for enzyme production by recombinant DNA technology. This was due to its promising trait that is suitable for genetically modify and the obtainability of a difference *E. coli* strains. But, the main downside when utilizing *E. coli* as a host is bacterial cell lysis due to the pressure build-up through overproduction of the expressed recombinant enzyme in the periplasmic space. Thus, the utilization of the immobilization in targeting the recombinant enzyme expression in the culture medium, presents substantial preferences over cytoplasmic excretion. Immobilization process can be applied to optimize the operational performance system of cell for industrial applications which leads to the development of economically and ecologically available enzyme such as xylanase. In this study, the effects of graphene oxide (GO) on xylanase excretion and β -galactosidase activity of immobilized *E. coli* was examined. The experiments were performed under the optimized RSM conditions (20 °C, 0.5 mM, IPTG and pH 7) using shake flask cultivation followed by bioreactor conditions (20 °C, 0.5 mM, IPTG and pH 7) in batch fermentation. The immobilized cells onto GO using shake flask cultivation exhibit about 3.8-fold increase in xylanase production (0.4821 U/ml) or equal to 73.7% higher compared to free cells (0.1264 U/ml). The *E. coli* were also demonstrated 15.3% reduction of cell lysis (0.0299 U/ml of β -galactosidase activity) compared to free cells (0.1273 U/ml of β -galactosidase activity) and 72.6% increase in plasmid stability compared to free cells. The xylanase concentration was 0.482 U/ml, representing 99.4% of the predicted value (0.485 U/ml) and 1.71-fold higher than the value before optimization process (0.283 U/ml). The occurrence of cell lysis demonstrated 99.3% reduction and increased in the plasmid stability up to 87% under the optimized RSM condition compared to free cells. The stirrer tank bioreactor also showed 5% higher in xylanase excretion (0.506 U/ml) with 33% reduction of β -galactosidase activity (0.06 U/ml) compared to shake flask cultivation (0.09 U/ml) after 24h. While from the Liquid Chromatography Mass Spectrometry (LCMS) analysis showed that the xylooligosaccharides was found to be hydrolyzed by an endoxylanase to produce xylopentose sugar. Hence, this study demonstrated that the immobilization of *E. coli* on GO potentially to be valuable for the excretion of recombinant proteins in *E. coli* with high cell viability.

ABSTRAK

Escherichia coli adalah sistem hos utama untuk penghasilan enzim menggunakan teknologi rekombinan DNA disebabkan ciri-ciri yang memberangsangkan untuk pengubahsuaian genetic juga mudah untuk didapati. Namun, kelemahan utama apabila menggunakan *E. coli* sebagai hos adalah pemecahan sel bakteria; akibat tekanan yang terbina melalui pengeluaran enzim secara berlebihan dalam ruang periplasma. Penggunaan pendekatan imobilisasi dalam mensasarkan penghasilan enzim rekombinan dalam medium kultur, menjadi pilihan utama berbanding perkumuhan sitoplasma. Proses imobilisasi boleh digunakan untuk sel bagi mengoptimumkan sistem operasi untuk aplikasi industri yang membawa kepada pembangunan biomangkin ekonomi dan ekologi enzim seperti xylanase. Dalam projek ini, kesan oksida grafit (GO) pada rembesan xylanase dan aktiviti β -galactosidase dari *E. coli* telah diamati. Kajian ini telah dilakukan di bawah optimum kondisi RSM (20 °C, 0.5 mM IPTG dan pH 7) menggunakan kaedah kelalang bergoncang diikuti oleh fermentasi dalam bioreactor (20 °C, 0.5 mM IPTG dan pH 7) dengan menggunakan kultur *batch*. Sel imobilisasi menggunakan nanopartikel GO mempamerkan peningkatan 3.8 kali ganda dalam pengeluaran xylanase (0.4821 U/ml) atau bersamaan dengan 73.7% lebih tinggi berbanding sel bebas (0.1264 U/ml). Imobilisasi sel juga menunjukkan pengurangan kadar sel pecah sebanyak 15.3% (0.0299 U/ml daripada aktiviti β -galactosidase) berbanding sel bebas (0.1273 U / ml aktiviti β -galactosidase) dan peningkatan dalam kestabilan plasmid sebanyak 72.6% berbanding sel bebas. Xylanase mempamerkan kepekatan sebanyak 0.482 U/ml, mewakili 99.4% daripada nilai yang diramalkan (0.485 U/ml) dan 1.71 kali ganda lebih tinggi daripada nilai sebelum proses pengoptimuman (0.283 U/ml). Kadar lisis sel menunjukkan pengurangan 99.3% dan peningkatan dalam kestabilan plasmid sehingga 87% di bawah kondisi optimum RSM berbanding sel bebas. Agitasi bioreaktor juga menunjukkan peningkatan perembesan xylanase sebanyak 5% lebih tinggi dalam pengeluaran xylanase (0.506 U/ml) dengan pengurangan aktiviti β -galactosidase aktiviti sebanyak 33% (0.06 U/ml) berbanding kelalang kon; (0.09 U/ml) selepas 24 jam. Manakala melalui Kromatografi Cair-Spektrometri Massa (LCMS) menunjukkan bahawa xylooligosakarida didapati dihidrolisis oleh endoxylanase untuk menghasilkan gula xylopentaosa. Jadi, kajian ini membuktikan imobilisasi *E. coli* pada GO berguna untuk perembesan rekombinan protin dalam *E. coli* dengan kestabilan sel yang tinggi.

ABSTRACT

Escherichia coli is the preference host system for enzyme production by recombinant DNA technology. This was due to its promising trait that is suitable for genetically modify and the obtainability of a difference *E. coli* strains. But, the main downside when utilizing *E. coli* as a host is bacterial cell lysis due to the pressure build-up through overproduction of the expressed recombinant enzyme in the periplasmic space. Thus, the utilization of the immobilization in targeting the recombinant enzyme expression in the culture medium, presents substantial preferences over cytoplasmic excretion. Immobilization process can be applied to optimize the operational performance system of cell for industrial applications which leads to the development of economically and ecologically available enzyme such as xylanase. In this study, the effects of graphene oxide (GO) on xylanase excretion and β -galactosidase activity of immobilized *E. coli* was examined. The experiments were performed under the optimized RSM conditions (20 °C, 0.5 mM, IPTG and pH 7) using shake flask cultivation followed by bioreactor conditions (20 °C, 0.5 mM, IPTG and pH 7) in batch fermentation. The immobilized cells onto GO using shake flask cultivation exhibit about 3.8-fold increase in xylanase production (0.4821 U/ml) or equal to 73.7% higher compared to free cells (0.1264 U/ml). The *E. coli* were also demonstrated 15.3% reduction of cell lysis (0.0299 U/ml of β -galactosidase activity) compared to free cells (0.1273 U/ml of β -galactosidase activity) and 72.6% increase in plasmid stability compared to free cells. The xylanase concentration was 0.482 U/ml, representing 99.4% of the predicted value (0.485 U/ml) and 1.71-fold higher than the value before optimization process (0.283 U/ml). The occurrence of cell lysis demonstrated 99.3% reduction and increased in the plasmid stability up to 87% under the optimized RSM condition compared to free cells. The stirrer tank bioreactor also showed 5% higher in xylanase excretion (0.506 U/ml) with 33% reduction of β -galactosidase activity (0.06 U/ml) compared to shake flask cultivation (0.09 U/ml) after 24h. While from the Liquid Chromatography Mass Spectrometry (LCMS) analysis showed that the xylooligosaccharides was found to be hydrolyzed by an endoxylanase to produce xylopentose sugar. Hence, this study demonstrated that the immobilization of *E. coli* on GO potentially to be valuable for the excretion of recombinant proteins in *E. coli* with high cell viability.

ABSTRAK

Escherichia coli adalah sistem hos utama untuk penghasilan enzim menggunakan teknologi rekombinan DNA disebabkan ciri-ciri yang memberangsangkan untuk pengubahsuaian genetic juga mudah untuk didapati. Namun, kelemahan utama apabila menggunakan *E. coli* sebagai hos adalah pemecahan sel bakteria; akibat tekanan yang terbina melalui pengeluaran enzim secara berlebihan dalam ruang periplasma. Penggunaan pendekatan imobilisasi dalam mensasarkan penghasilan enzim rekombinan dalam medium kultur, menjadi pilihan utama berbanding perkumuhan sitoplasma. Proses imobilisasi boleh digunakan untuk sel bagi mengoptimumkan sistem operasi untuk aplikasi industri yang membawa kepada pembangunan biomangkin ekonomi dan ekologi enzim seperti xylanase. Dalam projek ini, kesan oksida grafit (GO) pada rembesan xylanase dan aktiviti β -galactosidase dari *E. coli* telah diamati. Kajian ini telah dilakukan di bawah optimum kondisi RSM (20 °C, 0.5 mM IPTG dan pH 7) menggunakan kaedah kelalang bergoncang diikuti oleh fermentasi dalam bioreactor (20 °C, 0.5 mM IPTG dan pH 7) dengan menggunakan kultur *batch*. Sel imobilisasi menggunakan nanopartikel GO mempamerkan peningkatan 3.8 kali ganda dalam pengeluaran xylanase (0.4821 U/ml) atau bersamaan dengan 73.7% lebih tinggi berbanding sel bebas (0.1264 U/ml). Imobilisasi sel juga menunjukkan pengurangan kadar sel pecah sebanyak 15.3% (0.0299 U/ml daripada aktiviti β -galactosidase) berbanding sel bebas (0.1273 U / ml aktiviti β -galactosidase) dan peningkatan dalam kestabilan plasmid sebanyak 72.6% berbanding sel bebas. Xylanase mempamerkan kepekatan sebanyak 0.482 U/ml, mewakili 99.4% daripada nilai yang diramalkan (0.485 U/ml) dan 1.71 kali ganda lebih tinggi daripada nilai sebelum proses pengoptimuman (0.283 U/ml). Kadar lisis sel menunjukkan pengurangan 99.3% dan peningkatan dalam kestabilan plasmid sehingga 87% di bawah kondisi optimum RSM berbanding sel bebas. Agitasi bioreaktor juga menunjukkan peningkatan perembesan xylanase sebanyak 5% lebih tinggi dalam pengeluaran xylanase (0.506 U/ml) dengan pengurangan aktiviti β -galactosidase aktiviti sebanyak 33% (0.06 U/ml) berbanding kelalang kon; (0.09 U/ml) selepas 24 jam. Manakala melalui Kromatografi Cair-Spektrometri Massa (LCMS) menunjukkan bahawa xylooligosakarida didapati dihidrolisis oleh endoxylanase untuk menghasilkan gula xylopentaosa. Jadi, kajian ini membuktikan imobilisasi *E. coli* pada GO berguna untuk perembesan rekombinan protin dalam *E. coli* dengan kestabilan sel yang tinggi.

TABLE OF CONTENTS

TITLE	i
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
TABLE OF CONTENTS	vii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF SYMBOLS	xvii
LIST OF ABBREVIATION	xvii
LIST OF APPENDICES	xviii
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Problem statement	3
1.3 Objectives of the study	5
1.4 Scopes of the study	5
1.5 Rationale and significance of the research	6

CHAPTER 2 LITERATURE REVIEW 7

2.1	Background	7
2.2	Engineered Thermostable xylanase	8
2.3	System for producing recombinant proteins	9
2.3.1	<i>Escherichia coli</i> expression system	9
2.3.2	Yeast	10
2.3.3	Filamentous Fungi	11
2.4	Whole Cell Immobilization	13
2.5	Methods of Immobilization	14
2.5.1	Entrapment	14
2.5.2	Adsorption	15
2.5.3	Covalent Binding	15
2.6	Choice of Matrices for Immobilization	16
2.6.1	Nanoparticle as Immobilization Matrices	17
2.7	Factors Affecting the Cell Immobilization System	19
2.7.1	Cell Matrix Interaction	19
2.7.2	Medium	20
2.7.3	pH	21
2.7.4	Temperature	23
2.7.5	Agitation Rate	24
2.7.6	Inducer Concentration	25
2.8	Bioconversion of xylan into value-added biomolecules using xylanase	26

CHAPTER 3 METHODOLOGY**29**

3.1	Research design for the improvement of engineered thermostable xylanase excretion by immobilized recombinant <i>E. coli</i>	29
3.2	<i>E. coli</i> strains and plasmid selection	30
3.3	Chemicals and reagents	30
3.4	Preparation of bacterial glycerol stock	30
3.5	Cell immobilization	30
3.6	Expression study on engineered thermostable xylanase excretion by immobilized recombinant <i>E. coli</i>	31
3.6.1	Screening of the cultural conditions using one factor at one time method (OFAT)	31
3.7	Experimental design on engineered thermostable Xylanase excretion by the immobilized recombinant <i>E.coli</i>	36
3.7.1	Screening of the cultural conditions using full factorial design (FFD)	36
3.7.2	Optimization of the cultural conditions by Response Surface Methodology (RSM)	37
3.8	Reusability of immobilized cells	38
3.9	Batch cultivation in stirred-tank bioreactor	39
3.10	Preparation of enzymes	39
3.11	Analysis Procedures	39
3.11.1	Xylanase activity	40
3.11.2	Beta-galactosidase activity (cell lysis)	40
3.11.3	Plasmid stability	40

3.11.4	Cell density	41
3.11.5	Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE)	41
3.11.6	Field Emission Scanning Electron Microscope (FESEM)	42
3.11.7	Xylan Reaction Process	42
3.11.8	Solid Phase Extraction (SPE)	42
3.11.9	Liquid Chromatography Mass Spectrometry (LCMS)	43

CHAPTER 4 RESULTS AND DISCUSSION

4.1	Screening of cultivation conditions on xylanase excretion and cell viability of immobilized cell using OFAT	44
4.1.1	Effect of nanoparticles on xylanase excretion and cell viability of immobilized <i>E. coli</i>	44
4.1.2	Effect of expression medium on xylanase excretion and cell viability of immobilized <i>E. coli</i>	46
4.1.3	Effect of inducer concentration on xylanase excretion and cell viability of immobilized <i>E. coli</i>	49
4.1.4	Effect of post induction temperature on xylanase excretion and cell viability of immobilized <i>E. coli</i>	51
4.1.5	Effect of induction time on xylanase excretion and cell viability of immobilized <i>E. coli</i>	53
4.1.6	Effect of agitation rate on xylanase excretion and cell viability of immobilized <i>E. coli</i>	55

4.1.7	Effect of pH selection on xylanase excretion and cell viability of immobilized <i>E. coli</i>	58
4.1.8	Effect of inoculum size on xylanase excretion and cell viability of immobilized <i>E. coli</i>	60
4.2	Screening of cultural conditions on xylanase excretion and cell viability of immobilized <i>E. coli</i> using design of experiment (full factorial design)	61
4.3	Optimization of cultural conditions on engineered thermostable xylanase excretion and cell viability of immobilized <i>E. coli</i> using response surface methodology (central composite design)	66
4.4	Xylanase excretion, β -galactosidase activity, dry cell weight and plasmid stability between free cell, immobilized cell using OFAT, RSM and bioreactor condition	74
4.5	Reusability of immobilized recombinant <i>E. coli</i>	79
4.6	Characterization of Immobilized <i>Escherichia coli</i>	83
4.6.1	FESEM characterization	83
4.6.2	Performance and kinetic parameter values of recombinant xylanase by using the different type of cultivation conditions (Free Cell, OFAT, RSM & Bioreactor).	85
4.6.3	Liquid chromatography mass spectrometry (LCMS) analysis of xylooligosaccharides production	87
CHAPTER 5 CONCLUSION AND RECOMMENDATION		89
5.1	Conclusion	89
5.2	Recommendation	90
REFERENCES		92
APPENDICES		107
VITA		

LIST OF TABLES

3.1	The list of independent variables and parameter value of OFAT	32
3.2	The OFAT cultivation condition for improvement of xylanase production	32
3.3	The list of different expression media and formulation	34
3.4	Independent variables and the levels of the screening design	36
3.5	Centre point details for each parameter used in statistical design	37
3.6	Experimental design of the central composite design	37
4.1	Experimental design and results of full factorial design.	62
4.2	Analysis of variance for selected factorial model of xylanase excretion	64
4.3	Analysis of variance for selected factorial model of β -galactosidase activity	64
4.4	Statistical analysis for xylanase excretion and β -galactosidase activity	65
4.5	Experimental design and results of central composite design.	67
4.6	Analysis of variance for response surface quadratic model for xylanase excretion.	68
4.7	Analysis of variance for response surface quadratic model for β -galactosidase activity.	69
4.8	Summary of the optimized cultural conditions for the xylanase excretion and β -galactosidase activity	73

4.9	Performance and kinetic parameter values of free and immobilized recombinant E. coli	86
F1	Diagnostics case statistics of the xylanase activity (FFD)	112
F2	Diagnostics case statistics of the β -galactosidase activity (FFD)	113
G1	Diagnostics case statistics of the xylanase activity (RSM)	115
G2	Diagnostics case statistics of the β -galactosidase activity (RSM)	115



LIST OF FIGURES

2.1	Methods of immobilization	14
3.1	Research design overview for improvement of xylanase production	29
3.2	The workflow chart of Solid Phase Extraction steps	43
4.1	The effect of nanoparticle types on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	45
4.2	The effect of expression medium on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	48
4.3	The effect of IPTG concentrations on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	49
4.4	The effect of post induction temperature on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	52
4.5	The effect of induction time on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	54
4.6	The growth profile of Recombinant <i>Escherichia coli</i> . The middle log phase was equivalent to the 5 hr	54
4.7	The effect of agitation rate on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	57
4.8	The effect of pH on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	58

4.9	The effect of inoculum size on xylanase excretion and β -galactosidase activity of immobilized <i>E. coli</i> .	60
4.10	Response surface plot of xylanase excretion: temperature vs. induction time. The xylanase excretion of immobilized cell was measured after 24 hr induction time.	71
4.11	Response surface plot of xylanase excretion: IPTG concentration vs. induction time. The xylanase excretion of immobilized cell was measured after 24 hr induction time.	72
4.12	Response surface plot of β -galactosidase activity: induction time vs. IPTG concentration. The β -galactosidase activity of immobilized cell was measured after 24 hr induction time.	72
4.13	Xylanase excretion of free cell, immobilized cell using OFAT condition, immobilized cell using RSM condition and bioreactor.	75
4.14	β -galactosidase activity of free cell, immobilized cell using OFAT condition, immobilized cell using RSM condition and bioreactor.	76
4.15	Dry cell weight of free cell, immobilized cell using OFAT condition, immobilized cell using RSM condition and bioreactor.	77
4.16	Plasmid stability of free cell, immobilized cell using OFAT condition, immobilized cell using RSM condition and bioreactor.	79
4.17	Xylanase excretion between immobilized cell (onto non-treated graphene oxide) and free cell by repeated fermentation	80
4.18	β -galactosidase activity of immobilized cell (onto non-treated graphene oxide) and free cell by repeated fermentation	81

4.19	Plasmid stability of immobilized cell (onto non-treated graphene oxide) and free cell by repeated fermentation	82
4.20	Dry cell weight of immobilized cell (onto non-treated graphene oxide) and free cell by repeated fermentation	83
4.21	FESEM image of graphene oxide at 30 k x magnification.	84
4.22	FESEM image of immobilized recombinant Escherichia coli cells at 30k x magnification	84
4.23	LCMS analysis for the quantification of xylooligosaccharides production using the waters ACQUITY UPLC BEH Amide column (100 mm x 2.1 mm x 1.7 μ m).	88
D1	Xylose standard curve for the calculation of xylanase assay	110
E1	ONP standard curve for the calculation of B-galactosidase activity	111



LIST OF SYMBOLS AND ABBREVIATIONS

g	-	Gram
h	-	Hour
l	-	Liter
min	-	Minute
ml	-	Milliliter
sec	-	Second
rpm	-	Revolution per minutes
<i>E. coli</i>	-	<i>Escherichia coli</i>
Sp.	-	Species
CCD	-	Central Composite Design
DO	-	Dissolved Oxygen
FFD	-	Full Factorial Design
CNT	-	Carbon Nanotube
LB	-	Luria Bertani
GO	-	Graphene Oxide
ONPG	-	<i>Ortho</i> -nitrophenyl- β -galactoside
RSM	-	Response Surface Methodology
ANOVA	-	Analysis of Variance
DNA	-	Deoxyribonucleic Acid
OFAT	-	One Factor at One Time
IPTG	-	Isopropyl β -D-1-thiogalactopyranoside
LCMS	-	Liquid Chromatography Mass Spectrometry
FESEM	-	Field Emission Scanning Electron Microscopy

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Medium Preparation	107
B	Buffer Preparation	108
C	Antibiotic and Inducer	109
D	Standard curve for xylanase assay	110
E	Standard curve for B-galactosidase activity	111
F	Results for screening process of cultural conditions using FFD	112
G	Results for optimization process of cultural conditions using RSM	115
H	Publications and awards	116

CHAPTER 1

INTRODUCTION

1.1 Introduction

Recombinant DNA technology has facilitated researchers to bring an extensive amount of various proteins and enzymes in microbes that were hard to gain in quantity or relatively high-priced. The foreign genes expression has been recorded in a range of microorganisms such as fungi and bacteria by cloning the desired gene into the plasmid vector under the regulation of an inducible promoter such as isopropyl β -D-1-thiogalactopyranoside (Kuniechick *et al.*, 2009). The expression of recombinant enzyme is recognized as heterologous protein; and the most frequently utilized expression systems for heterologous gene production of both prokaryotic and eukaryotic origins is microbial expression system (Fakruddin *et al.*, 2013).

The usage of plasmid as a vector for production of recombinant enzyme or protein is vital to produce high beneficial enzymes, in adequate supply for research and industrial prospect (Joseph *et al.*, 2015). However, in commercial production of recombinant microorganisms, the plasmid instability has become one of the major issues during this process. The plasmid instability is an inclination of the transformed cells to lose their engineered characteristics by alteration or loss of plasmid. As a consequence, it may impact in a critical loss in targeting recombinant protein throughput and the most often utilized bacterial expression system to produce the heterologous protein is *E. coli* (Fakruddin *et al.*, 2013; Kuniechick *et al.*, 2009).

Escherichia coli is Generally Recognized as Safe (GRAS) organism and verified to be a cost-effective host cell line for proteins and enzymes production by recombinant DNA technology, due to its promising traits such as fast and high cell growth, easy to cultivate, low cost, suitable and simple for genetically modify and the obtainability of a different *E. coli* strains (Rosano & Ceccarelli, 2014). However, in the process, tons of

issues can possibly happen, such as inactivation of the protein enzyme, formation of the inclusion body (IB) and unfavorable cell growth rate are certain complications that frequently occurred (Rosano & Ceccarelli, 2014; Birkholtz *et al.*, 2008).

Throughout the expression of heterologous protein, the composition of insoluble aggregates or lower production might result from the dissimilarities in identical codon usage of expression host. Other than that, the protein overexpression in *E. coli* could also lead to insoluble protein synthesis in the form of inclusion bodies, caused by metabolic burden. When a foreign DNA is brought into the bacterial strain, the new synthesis recombinant polypeptide will be expressed in the microenvironments of *E. coli*. This expression might vary from the initial condition relating to osmolarity, cofactors, folding mechanisms, pH, redox potential and both spatial and temporally control of gene expression is disoriented (Rosano & Ceccarelli, 2014). Hence, the desired protein production involves bioprocess media optimization for targeting the high cell density culture. By determining the optimum cultivation condition such as medium type for microorganism can reduce the metabolic stress exposed to the host cells (Angov *et al.*, 2008; Birkholtz *et al.*, 2008).

Thus, it is worthy to discover the optimum medium for cultivation system such as by providing the enriched culture medium for higher soluble proteins expression. There are several types of media have been used for cultivation such as, Terrific Broth media, salt media and Luria-Bertani broth (Baser *et al.*, 2010). While, the Luria-Bertani broth is the most regularly utilized medium culture for *E. coli* as it contains high nutrients, measures simple preparation and optimum osmolarity for the early exponential growth phase. All these aspects make it sufficient for enzyme production and recompense for the statement claiming it is not the best preference for attaining high cell density cultures (Sezonov & Ari, 2007).

On another point of view, to improve the immobilization process for better operational performance system of recombinant *E. coli* and with the development of nanotechnology, it is now possible for nanoparticles to be used as immobilization matrix for a wide range of industrial applications such as production of recombinant protein by immobilized bacteria. This is due to nanoparticle's specific physico-chemical properties such as its nano scale size, chemical composition and surface area that make it suitable material to fix diffusion problems when the process is dealing with macromolecular substrates. Moreover, a large surface-to-volume proportion presented by nanomaterial resulted in the higher concentration of the immobilized cells compared to those gave by other materials (Ahmad & Sardar, 2015; Saifuddin *et al.*, 2016). Thus, nanoparticles such

as graphene oxide and carbon nanotube are key modules in the future market of advance innovation that greatly impact the material's mechanical properties as well as biocompatible environment or surrounding area for immobilization method (Ahmad & Sardar, 2015).

Concurrently, the substrate utilized for this research; polysaccharides of xylan are generally correlated with lignin as well as cellulose, which act as an essential structural support to build up the plant cell walls. The degradation of hemicellulose constituent, predominantly comprised of xylan, necessitates effective xylanolytic enzymatic systems involving certain enzymes such as β -xylosidase, endo- β -1,4-xylanase, α -glucuronidase, α -L-arabinofuranosidase, p-coumaric acid esterase, acetyl xylan esterase and ferulic acid esterase. Recently, there has been a rising attention in xylanase enzyme fermentations concerning their wide-ranging applications in a number of industries, particularly in the pulp and paper commerce (Rusli *et al.*, 2009; Basar *et al.*, 2010).

Therefore, xylanases enzymes accountable for the hydrolysis of xylan to xylooligosaccharides (XOS) and have drawn intentness by virtue of their prospective in numerous industrial progresses. Moreover, xylose is the primary constituent of the hemicellulose xylan and the secondary most sufficient plant biomass derivative sugar (Salamanca *et al.*, 2014). Xylanase in combination with other enzymes, act efficiently to degenerate xylan to sugar constituents which can be produce naturally by wild type filamentous fungi species such as *Trichoderma sp.* and *Aspergillus sp.* Xylanase can be applied in wide-ranging utilization for industrial scales such as to enhance the quality of bread, improve the digestibility of ruminant feeds and as a pre-bleaching of kraft pulp (Farliahati *et al.*, 2010).

In this study, the graphene oxide was chosen as a matrix for recombinant *E. coli* to improve the engineered thermostable xylanase excretion with fewer occurrences of cell lysis. Furthermore, a thorough search for relevant literatures did not yield any related article and to the extent of our knowledge, there have been no studies on excretion of engineered thermostable xylanase by recombinant *E. coli* that have been immobilized through an adsorption of the cells using graphene oxide.

1.2 Problem statement

The establishment of heterologous protein by employing recombinant DNA innovation has drawn a great interest in the industrial process of biotechnology and the manufacturing level (Man *et al.*, 2015; Mergulhão *et al.*, 2005). However, despite

impressive progress in employing the *E. coli* cell factory for excretory production of recombinant enzymes, the production of expressed protein has encountered with the certain operative obstacle during the process (Yoon *et al.*, 2010).

Firstly, the complications in recovering excretion of correctly folded proteins; as the overexpression of proteins usually occur in the inclusion bodies (IB) form from the active proteins that could only be recovered by a complex and expensive process (Choi & Lee, 2004). Secondly, the excretion yield of recombinant protein was usually untraceable or low-set; mainly because of interactions within the hydrophobic surface regions of protein causing aggregates (Yoon *et al.*, 2010). Thirdly, the autolysis of cell due to overexpression of recombinant protein in host cell (Choi & Lee, 2004).

Accordingly, assortment techniques have been utilized and established to resolve these issues; for instance, lowering the concentration of inducer by controlling expression level using various promoters, excretion of recombinant proteins into the culture medium or periplasmic space, use of various host strains and certain recombinant proteins to induce modulation of chaperones and a class of proteins that assist protein folding (Wyre & Overton, 2014; Choi & Lee, 2004). The other commonly suitable method to maximize recombinant protein solubility and folding is by reducing the imposed stress on bacteria and lowering the rates of recombinant protein excretion by controlling the growth temperature. This is to enable the translation and folding process occurs more gradually in recombinant protein (Wyre & Overton, 2014).

Moreover, the declined yields of a target protein through the gene expression of *E. coli* are regularly caused by the metabolic stress and consequence in the reduced cell growth and redirection of cellular metabolic process. In certain cases, it has been described to occur caused by the addition of isopropyl β -D-1-thiogalactopyranoside (IPTG) inducer. Specifically, once induction is initiated when a certain critical cell mass is obtained. This is probably caused by the deficiency of essential metabolites during process (Ramchuran *et al.*, 2005). Furthermore, the cultural conditions and environmental factors such as temperature have the main influence on the total protein expression and cell metabolism (Morowvat *et al.*, 2015).

Thus, in order to improve the system performance of the recombinant *E. coli* for recombinant protein excretion, immobilization of cell was used. Cell immobilization could reduce the plasmid instability and cell lysis issues, improving the productiveness of the recombinant cells, low sensitivity to pH and temperature when compared with free cells. Thus, will lowering the possibility of contamination, decreasing the fermentation time, advancing the rate of substrate uptake, and resultant in estimated concentration of

volatile component for product. Meanwhile, studies on finding the convenient method to improve the xylanase enzyme production that being recognized as high potential enzyme for wide range application of industrial use such as pulp and paper industry is still deliberately studied by researchers. Consequently, this study demonstrates that graphene oxide is a convenient immobilization matrix for recombinant *E. coli* due to its unique features as well as special characteristics to enhance the immobilization cell efficiency and excrete high xylanase expression.

1.3 Objectives of the study

1. To improve the engineered thermostable xylanase excretion of immobilized recombinant *E. coli* onto graphene oxide in comparison with free cell.
2. To determine the cell viability with high plasmid stability of immobilized recombinant *E. coli* onto graphene oxide.

1.4 Scopes of the study

1. Screening the effect of the cultural conditions (nanomatrix quantity, expression medium, IPTG, post induction temperature, post induction time, agitation rate, inoculum size and pH) on improvement of xylanase excretion and cell viability of the immobilized cell using one factor at one-time method (OFAT).
2. Screening the effect of significant factors of cultural conditions (IPTG, post induction temperature, post induction time, agitation rate, inoculum size and pH) on the immobilized cell for improvement of xylanase excretion and cell viability using Design-Expert Software; full factorial design (FFD).
3. Optimization of the cultural conditions (IPTG, post induction temperature and post induction time) on the excretion of recombinant xylanase and reduction of cell lysis using Design-Expert Software; Response Surface Methodology (RSM).
4. Production of high level xylanase excretion of immobilized cell in the 5 L bioreactor by using optimum RSM conditions.
5. Determination of immobilized cell's enzymatic production of xylooligosaccharides (XOS) from xylan using Liquid Chromatography-Mass Spectrometry (LCMS).

REFERENCES

- Adhikari, P. D., Jeon, S., & Cha, M. (2014). Immobilization of carbon nanotubes on functionalized graphene film grown by chemical vapor deposition and characterization of the hybrid material, *Science and Technology of Advanced Materials*, 15(1). doi:10.1088/1468-6996/15/1/015007
- Affney, M. G., Sean, D. O., & Richard, M. U. (2009). Optimization of xylanase production by *Thermomyces lanuginosus* in solid state fermentation. *Bioscience, Biotechnology, and Biochemistry*, 73(12), 2640–44. doi: 10.1271/bbb.90493
- Afzali, N., Bahareh, R. Z., & Nahidhagh, N. (2016). Optimization of *Trichoderma reesei* medium for increasing xylanase enzyme production. *Advances in BioResearch* 7, 94–99. doi: 10.15515/abr.0976-4585.7.3.9499
- Ashikin, N., A., L., Wahab, M., K., Illias, R. & Fuzi, S., F., Z. (2017). “Comparison of Thermostable Xylanase Production by *Escherichia Coli* Immobilised onto Different Nanoparticles.” *The Italian Association of Chemical Engineering* 56: 1825–1830. doi: 10.3303/CET1756305
- Ahmad, R., & Sardar, M. (2015). Biochemistry & analytical biochemistry enzyme immobilization: an overview on nanoparticles as immobilization matrix. *Biochemistry & Analytical Biochemistry*, 4(2). doi:10.4172/2161-1009.1000178
- Akhavan, O., Ghaderi, E., & Esfandiar, A. (2015). Wrapping bacteria by graphene nanosheets for isolation from environment, reactivation by sonication, and inactivation by near - infrared irradiation, 1–17. doi:10.1021/jp200686k
- Ali, M. B., Irshad, M., Anwar, Z. & Zafar, M. (2016). Screening and statistical optimization of physiochemical parameters for the production of xylanases from agro-industrial wastes. *Advances in Enzyme Research*, 4, 20–33. doi: 10.4236/aer.2016.41003
- Ammar, A., Al, A. M., Alali, M., & Karim, A. (2016). Influence of graphene oxide on mechanical, morphological, barrier, and electrical properties of polymer membranes. *Arabian Journal of Chemistry*, 9(2), 274–286.

- Anazawa, H., & Sakayu, S. (2014). Microbial production: from genome design to cell engineering. In *Springer Science & Business Media*, 145. doi: 10.1007/978-4-431-54607-8
- Angov, E., Hillier, C. J., Kincaid, R. L., & Lyon, J. A. (2008). Heterologous protein expression is enhanced by harmonizing the codon usage frequencies of the target gene with those of the expression host, *PLoS One*, 3(5), 1–10.
- Arulraj, P., Kandasamy, Bhuvaneswari, A. & Venkatanarayanan, R. (2015). “Design, Development and Comparison of Fermentative Production of Ethanol by Free and Immobilized *Saccharomyces Cerevisiae* Under Stationary Culture.” *Journal of Pharmacy and Pharmaceutical Sciences* 4(9): 950–58.
- Asadollahi, N., Yavari, R., & Ghanadzadeh, H. (2015). Preparation, characterization and analytical application of stannic molybdophosphate immobilized on multiwalled carbon nanotubes as a new adsorbent for the removal of strontium from wastewater, *Journal of Radioanalytical and Nuclear Chemistry*, 303(3), 1–26.
- Assamoi, A. A., Delvigne, F., Aldric, J. & Destain, J. (2017). Improvement of Xylanase production by *Penicillium canescens* 10-10c in solid-state fermentation. *Biotechnology, Agronomy, Society and Environment*, 12, 1–9.
- Aoyagi, Hideki, Yoichi Katakura, and Akio Iwasaki. (2016). “Production of Secretory Cutinase by Recombinant *Saccharomyces Cerevisiae* Protoplasts.” *SpringerPlus* 5(160): 4–9. doi: 10.1186/s40064-016-1806-4
- Babaeipour, V., Seyed, A. S., & Nader, M. (2015). Maximizing production of human interferon- γ in HCDC of recombinant *E. Coli*. *Iranian Journal Pharmaceutical Research*, 12(3), 563–72.
- Bairagi, S. & Rohtak, H. (2016). Optimization of Cellulase Enzyme from Vegetable Waste by Using *Trichoderma Atroviride* in Solid State Fermentation. *Journal of Environmental Science, Toxicology and Food Technology*, 10(5), 68–73. doi: 10.9790/2402-1005026873
- Barbotin, J. N. (2001) Plasmid stability in immobilized cells. In: Wijffels R.H. (eds) *Immobilized Cells*. Springer (Lab Manuals). Springer, Berlin, Heidelberg. doi: 10.1007/978-3-642-56891-6_21
- Basar, B., Shamzi, M. M., Rosfarizan, M., Puspaningsih, N. N. T., & Ariff, A. B. (2010). Enhanced production of thermophilic xylanase by recombinant *Escherichia coli* DH5 α through optimization of medium and dissolved oxygen level. *International Journal of Agriculture & Biology*, 12(3), 321–328.

- Battestin, V., & Macedo, G. A. (2007). Effects of temperature, pH and additives on the activity of tannase produced by *Paecilomyces variotii*, *Electronic Journal of Biotechnology*, 10(2). doi:10.2225/vol10-issue2-fulltext-9
- Behnam, S., Karimi, K., Khanahmadi, M. & Salimian, Z. (2016). Optimization of xylanase production by *Mucor indicus*, *Mucor hiemalis*, and *Rhizopus oryzae* through solid state fermentation. *Biological Journal of Microorganism*, 4(16),1–10.
- Bekatorou, A., Stavros, P., & Athanasios, M. (2016). “Cell Immobilization Technologies for Applications in Alcoholic Beverages.” In Taylor & Francis Group, LLC, 4 , 933–55.
- Beshay, U., El-Enshasy, H., Ismail, I. M. K., Moawad, H., & Abd-El-Ghany, S. (2011). β -glucanase productivity improvement via cell immobilization of recombinant *Escherichia coli* cells in different matrices. *Polish Journal of Microbiology*, 60(2), 133–138. Retrieved from <http://www.scopus.com/inward/record.url?eid=2-s2.0-80054903645&partnerID=tZOtx3y1>
- Bill, R. (2016). Playing catch-up with *Escherichia coli*: using yeast to increase success rates in recombinant protein production experiments. *Microbiology*, 5(85), 1–12.
- Birkholtz, L., Blatch, G., Coetzer, T. L., Hoppe, H. C., Human, E., Morris, E. J., & Louw, A. I. (2008). Heterologous expression of plasmodial proteins for structural studies and functional annotation, *Malaria Journal*, 7(197), 1–20. <http://doi.org/10.1186/1475-2875-7-197>
- Brena, B., González-Pombo, P., & Batista-Viera, F. (2013). Immobilization of enzymes: a literature survey. *Immobilization of Enzymes and Cells*, 1051, 15–31. doi:10.1007/978-1-62703-550-7_2
- Briand, L., Marcion, G., Kriznik, A. & Heydel, J. M. (2016). A self-inducible heterologous protein expression system in *Escherichia coli*. *Scientific Reports*, 6, 1–11. doi: 10.1038/srep33037
- Burlacu, A., Călina, P., C., & Florentina, I. R. (2016). “Microbial Xylanase : A Review.” scientific bulletin: 335–42.
- Canbolat, M. F., Gera, N., Tang, C., Monian, B., Rao, B. M., Pourdeyhimi, B., & Khan, S. A. (2013). Preservation of cell viability and protein conformation on immobilization within nanofibers via electrospinning functionalized yeast. *ACS Applied Materials and Interfaces*, 5(19), 9349–9354. doi:10.1021/am4022768
- Cavalieri, N., Guimaraes, D. A., Carvalho, S. D., & Nogueira, P. (2013). Effect of Xylanase from *Aspergillus niger* and *Aspergillus flavus* on pulp biobleaching and enzyme production using agroindustrial residues as substract. *Bioprocess and*

Biotechnology, 2–9. doi: 10.1186/2193

- Chakdar, H., Murugan, K., Kuppusamy P., & Arjun S. (2016). Bacterial xylanases: biology to biotechnology. *Biotechnology*, 150(6). doi: 10.1007/s13205-016- 0457-z
- Chang, X., Xu, B., Bai, Y., Luo, H., Ma, R., & Shi, P. (2017). Role of N-linked glycosylation in the enzymatic properties of a thermophilic GH 10 xylanase from *Aspergillus fumigatus* expressed in *Pichia pastoris*. *PLoS ONE* 12(2). e0171111. doi:10.1371/journal.pone.0171111
- Chantorn, S., Chanitchote, P., Phataraporn, K., & Srisuda, P. (2016). Suitable conditions for xylanases activities from *Bacillus* Sp. GA2 (1) and *Bacillus* Sp. GA1 (6) and their properties for agricultural residues hydrolysis. *Journal of Science and Technology*, 38(2), 177–82.
- Che, R., Fauzi, A., Fatimah, S. & Illias, R. (2016). “Effects of culture conditions of immobilized recombinant *Escherichia coli* on *Cyclodextrin Glucanotransferase* (CGTase) excretion and cell stability. *Process Biochemistry*, 51, 474–83. doi: 10.1016/j.procbio.2016.01.002
- Chen, S., Mårten, L., Robert C. R., & Swaine, L. C. (2017). Direct and convenient measurement of plasmid stability in lab and clinical isolates of *E. Coli*. *Scientific Reports*, 7, 1–11. doi: 10.1038/s41598-017-05219-x
- Chen, S., & Chiu, B. K. (2009). A fermentation strategy for production of recombinant protein subjected to plasmid instability. *Korean Journal of Chemical Engineering*, 25(5), 1110–1114.
- Choi, J. H., & Lee, S. Y. (2004). Secretory and extracellular production of proteins using *Escherichia coli*. *Applied Microbiology and Biotechnology*, 64(5), 625–635.
- Comlekcioglu, U., Gunes, M., Altun, H., Ekiz, D. O., & Aygan, A. (2017). Coexpression of rumen fungal xylanase and bifunctional cellulase genes in *Escherichia coli*. *Brazilian Archives of Biology and Technology*, 60, 1–10. doi: 10.1590/1678-4324-2017160462
- Das, B., Chakraborty, A., & Ghosh, S. (2011). Studies on the effect of pH and carbon sources on enzyme activities of some pectinolytic bacteria isolated from jute retting water. *Turkish Journal of Biology*, 35, 671–678. doi:10.3906/biy-1008-50
- Duarte, J. Rodrigues, J., Moran, P., & Valenca, G., (2013). Effect of immobilized cells in calcium alginate beads in alcoholic fermentation. *AMB Express*, 3(1), 31. doi: 10.1186/2191.
- Dogan, N. M., Tugba, S., Gulumser, A. D., & Naime, N. B. (2016). “Immobilization of *Lysinibacillus Fusiformis* B26 Cells in Different Matrices for Use in Turquoise Blue

- HFG Decolourization.” Archives of Environmental Protection 42(2): 92–99. doi: 10.1515/aep-2016-0013
- Eed, J. (2013). Factors affecting enzyme activity. *ESSAI*, 10(19). Available at: <http://dc.cod.edu/essai/vol10/iss1/19>
- Elakkiya, M., Thirumarimurugan, M., & Prabhakaran, D. (2016). “Methods of Cell Immobilization and Its Application.” International Journal of Innovative Research in Science, Engineering and Technology 5(4): 5429–33. doi: 10.15680/IJRSET.2016.0504175
- El-gendi, H., Azab, M. S., & Soliman, N. A. (2016). Application of plackett-burman design for optimization of alkaline protease and α -amylase production by the marine bacterium *Bacillus methylotrophicus* SCJ4. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 7(899), 899–909.
- Elnashar, M. (2010). Review article: immobilized molecules using biomaterials and nanobiotechnology. *Journal of Biomaterials and Nanobiotechnology*, 1(1), 61–77. doi:10.4236/jbnb.2010.11008
- Fakruddin, M., Mazumdar, R. M., Shahnewaj, K., Mannan, B., Chowdhury, A., & Hossain, N. (2013). Critical factors affecting the success of cloning, expression, and mass production of enzymes by recombinant *E. coli*, *ISRN Biotechnology*.
- Farliahati, M. R., Ramanan, R. N., & Mohamad, R. (2010). Enhanced production of xylanase by recombinant *Escherichia coli* DH5 α through optimization of medium composition using response surface methodology, *Annals of Microbiology*, 60(2), 279–285. <http://doi.org/10.1007/s13213-010-0038-z>
- Fatokun, E. N., Uchechukwu, U. N., Ademola, O. O., & Anthony, I. (2017). Optimization of process conditions for the production of holocellulase by a bacillus species isolated from Nahoon Beach sediments. *American Journal of Biochemistry and Biotechnology*, 13(2), 70–80. doi: 10.3844/ajbbsp.2017.70.80
- Franca, A., Cleiton J., Dangelly M., & Antonio M. (2015). Effect of pH and temperature on the production and activity *Schwanniomyces polymorphus* extracellular protease in fermentation medium. *African Journal of Microbiology Research*, 9(14), 1044–1052.
- Fuzi, S. F. Z. M., Farah, D. A. B., Abdul, M. A. M., & Rosli, M. I. (2011). Development and validation of a medium for recombinant endo- β -1, 4-xylanase production by *Kluyveromyces lactis*. *Annals of Microbiology*, 62(1). doi: 10.1007/s13213-011-0258-x

- Fuzi, S. F. Z. M., Jamaliah, M. J., Roshanida, A. R., & Rosli, M. I. (2014). Simplified feeding strategies for the fed-batch cultivation of *Kluyveromyces lactis* GG799 for enhanced recombinant xylanase production. *Bioprocess and Biosystems Engineering*, 37, 1887–1898. doi: 10.1007/s00449-014-1163-z
- Garlet, T. B., Weber, C. T., Klaic, R., Foletto, E. L., Jahn, S. L., Mazutti, M. A., & Kuhn, R. C. (2014). Carbon Nanotubes as Supports for Inulinase Immobilization, 14615–14624. doi:10.3390/molecules190914615
- Galih, P. R., & Esyanti, R. R. (2014). Effect of Immobilization on Cell Growth and Alkaloid Contents in Cell-Aggregate Culture of *Eurycoma longifolia* Jack, 2(2), 10–13.
- Gomes, A. F. S., Bruna, S. L. D., & Milla, A. B. (2016). Substrate and temperature effect on xylanase production by *Aspergillus fumigatus* using low cost agricultural wastes. *Bioscience Journal*, 32(4), 915–21.
- Groboillot, A., Boadi, D. K., Poncelet, D., & Neufeld, R. J. (1994). Immobilization of cells for application in the food industry. *Critical Reviews in Biotechnology* 14(2), 75–107.
- Guan, G., Zhao, P., Zhao, J., & Wang, M. (2016). Production and partial characterization of an alkaline xylanase from a novel fungus *Cladosporium oxysporum*. *BioMed Research International*, 1–8. doi: 10.1155/2016/4575024
- Gungormusler, Y., Mine, N. C., David, B. L., & Nuri, A. (2015). Cell immobilization for microbial production of 1,3-propanediol. *Biotechnology*, 1–13. doi: 10.3109/07388551.2014.992386
- Gurung, N., Ray, S., Bose, S., & Rai, V. (2013). A broader view: microbial enzymes and their relevance in industries, medicine, and beyond. *Biomed Research International*, 18.
- Hakimi, M. K., Anuar M., & Illias, R. M. (2016). Thermostability enhancement of xylanase *Aspergillus fumigatus* RT-1. *Journal of Molecular Catalysis B: Enzymatic*, 134(A), 154-163. doi.org/10.1016/j.molcatb.2016.09.020
- Hanson, M., Brorson, K., Moreira, A., & Rao, G. (2009). Comparisons of optically monitored small-scale stirred tank vessels to optically controlled disposable bag bioreactors. *Microbiology Cell Factories*, 7, 1–7.
- Ho, H. L. (2014). Effects of medium formulation and culture conditions on microbial xylanase production using agricultural extracts in submerged fermentation (SmF) and solid state fermentation (SsF): A review. *Biodiversity, Bioprospecting and Development*, 1(3), 1–13. doi: 10.4172/2376-0214.1000130

- Ho, H. L. (2016). Batch submerged fermentation in shake flask culture and bioreactor: influence of different agricultural residuals as the substrate on the optimization of xylanase production by *Bacillus subtilis* and *Aspergillus brasiliensis*. *Journal of Applied Biotechnology & Bioengineering*, 1(3), 1–9. doi: 10.15406/jabb.2016.01.00016
- Homaei, A. A., Reyhaneh, S., Fabio, V., & Roberto, S. (2013). Enzyme immobilization. *Journal of Chemical Biology*, 6(4), 185–205. doi: 10.1007/s12154
- Huang, R., Chao, Z., Hui, T., & Ruming, Z. (2013). Effect of culture conditions on production of xylanase by *Trichoderma reesei*. *Advanced Materials Research* 784, 856–860. doi:10.4028/www.scientific.net/AMR.781-784.856
- Hyun, J., Chang, K., & Yup, S. (2006). Production of recombinant proteins by high cell density culture of *Escherichia coli*. *Chemical Engineering Science*, 61, 876–885. doi:10.1016/j.ces.2005.03.031
- Irfan, M., Asghar, U. & Nadeem, M. (2016). Optimization of process parameters for xylanase production by *Bacillus* sp. in submerged fermentation. *Journal of Radiation Research and Applied Sciences*, 9, 139–147. doi: doi.org/10.1016/j.jrras.2015.10.008
- Ivanova, V., Petia, P., & Jordan, H. (2011). Application in the ethanol fermentation of immobilized yeast cells in matrix of alginate / magnetic nanoparticles, on chitosan-magnetite microparticles and cellulose-coated magnetic nanoparticles. *International Review of Chemical Engineering*, 3(2), 289–299.
- Jaqueline, M., Maciel, M., Castro, A., & Camarão, H. (2016). Industrial and biotechnological applications of ligninolytic enzymes of the basidiomycota: A review, *Electronic Journal of Biotechnology*, 13(6), 1–13. doi:10.2225/vol13
- Jiang, Y. M., Wang, Y. T. & Dong, Z. W. (2007). Effects of induction starting time and Ca^{2+} on expression of active penicillin G acylase in *Escherichia coli*. *Biotechnology Progress*, 23(5), 1031–1037. doi: 10.1021/bp070100
- Joanna, Ż., Urszula, G., & Wojcieszynska, D. (2016). Metabolic responses of bacterial cells to metabolic responses of bacterial cells. *Molecules*, 21(958), 1–15. doi: 10.3390/molecules21070958
- Joseph, B. C., Suthakaran P., Sankaranarayanan S., & Musti M. (2015). An overview of the parameters for recombinant protein expression in *Escherichia coli*. *Cell Science & Therapy*, 6(5), 1-7.
- Kanimozhi, K, & P K Nagalakshmi. (2014). “Original Research Article Xylanase Production from *Aspergillus Niger* by Solid State Fermentation Using Agricultural

- Waste as Substrate.” *Int. J. Curr. Microbiol. App. Sci.* 3(3): 437–46.
- Khusro, A., Kannan, B., Valan, M. & Agastian, P. (2016). Statistical optimization of thermo- alkali stable xylanase production from *Bacillus tequilensis* strain. *Electronic Journal of Biotechnology*, 22, 16–25. doi: 10.1016/j.ejbt.2016.04.002
- Kourkoutas, Y., Bekatorou, A., Banat, M., Marchant, R., & Koutinas, A. A. (2004). Immobilization technologies and support materials suitable in alcohol beverages production: a review. *Food Microbiology*, 21(4), 377–397. doi: 10.1016/j.fm.2003.10.005
- Krajewska, B., Teofil, J., & Jakub Z. (2014). Enzyme immobilization by adsorption: a review, *Adsorption*, 20,801–821. doi:10.1007/s10450-014-9623-y
- Krause, M., Ukkonen, K., Haataja, T., & Ruottinen, M. (2010). A novel fed-batch based cultivation method provides high cell-density and improves yield of soluble recombinant proteins in shaken cultures. *Microbiology Cell Factories*, 9(11), 1–11.
- Kuniechick, N., Munareto, R., Milech, T., Nunes, P., Chies, J., Basso, L. A., & Santos, S. (2009). Determination of the stability of recombinant plasmid containing the coding sequence of the human growth hormone in bioreactor, *XSalão de Iniciação Científica PUCRS*, 2829-2831.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680-685.
- Lee, H. K., Lee, J. K., Kim, M., & Lee C. J. (2010). Immobilization of lipase on single walled carbon nanotubes in ionic liquid, *Bulletin of the Korean Chemical Society* 31(3), 650–652. doi:10.5012/bkcs.2010.31.03.650
- Lieko, Carla, Della Torre, and Marina Kimiko Kadowaki. 2017. “Thermostable Xylanase from Thermophilic Fungi: Biochemical Properties and Industrial Applications.” *African Journal of Microbiology Research* 11(2): 28–37.
- Li, J., Jaitzig, J. & Lu, P. (2015). Scale-up bioprocess development for production of the antibiotic valinomycin in *Escherichia coli* based on consistent fed-batch cultivations. *Journal of Microbiology Cell*, 14(83), 1–13. doi: 10.1186/s12934-015-0272-y
- Li, S., Yang, X., Yang, S., Zhu, M., & Wang, X. (2016). Technology prospecting on enzymes: application, marketing and engineering, *Computational and Structural Biotechnology Journal*, 2(3), 1–14. doi:10.5936/csbj.201209017
- Li, Z., Manfred, N. & Ursula, R. (2014). The metabolic potential of *Escherichia coli* BL21 in defined and rich medium. *Microbial Cell Factories*, 13(45), 1–17.

- Ling, X. M., Xiang, Y. W., Ping M., & Yi, Y. (2016). Covalent immobilization of penicillin G acylase onto Fe₃O₄@chitosan magnetic nanoparticles. *Journal of Microbiology and Biotechnology*, 26(5), 829–836. doi: 10.4014/jmb.1511.11052
- Liu, L., Yang, H., Shin, H., & Li, J. (2013). How to achieve high-level expression of microbial enzymes strategies and perspectives. *Bioengineered*, 4(4), 212–223.
- Maier, R. M. (2009). Bacterial growth: review of basic microbiological concepts. *Environmental Microbiology*: 37–54.
- Mamvura, T. A., Iyuke, S. E., Sibanda, V., & Yah, C. S. (2012). Immobilization of yeast cells on carbon nanotubes. *South African Journal of Science*, 108(7-8), 90-97.
- Man, R. C., Ismail A. F., Ghazali N. F., Fuzi S. F. Z. M., & Illias R. M. (2015). Effects of the immobilization of recombinant *Escherichia coli* on cyclodextrin glucanotransferase (CGTase) excretion and cell viability. *Biochemical Engineering Journal*, 98, 91–98. doi:10.1016/j.bej.2015.02.013
- Man, R. C., Ismail, A. F., Ghazali, N. F., Fuzi, S. F. Z. M., & Illias, R. M. (2016). Effects of culture conditions of immobilized recombinant *Escherichia coli* on cyclodextrin glucanotransferase (CGTase) excretion and cell stability. *Process Biochemistry Journal*, 51(4), 474–483. doi:10.1016/j.procbio.2016.01.002
- Mergulhão F. J. M., Summers D. K., & Monteiro G. A. (2005). Recombinant protein secretion in *Escherichia coli*. *Biotechnology Advances*, 23(3), 177–202. doi:10.1016/j.biotechadv.2004.11.003
- Mmango, K., Ziyanda, K. O., & Uchechukwu, U. N. (2016). Optimization of cellulase and xylanase production by micrococcus species under submerged fermentation. *Journal Sustainability*, 8(11), 1–15. doi: 10.3390/su8111168
- Mohamad, N. R., Haziqah, N., Marzuki, C., Buang, N. A., Huyop, F., & Wahab, R. A. (2016). An overview of technologies for immobilization of enzymes and surface analysis techniques for immobilized enzymes, *Biotechnology & Biotechnological Equipment*, 29(2), 205–220. doi:10.1080/13102818.2015.1008192
- Morowvat, M. H., Babaeipour, V., Memari, H. R., & Vahidi, H. (2015). Optimization of fermentation conditions for recombinant human interferon beta production by *Escherichia coli* using the response surface methodology. *Jundishapur Journal of Microbiology*, 8(4), 1–11.
- Motta, F. L., Andrade, C. C. P., & Santana, M. H. A. (2013). A Review of xylanase production by the fermentation of xylan: classification, characterization and applications. In: Chandel AK, da Silva SS (eds) Sustainable degradation of lignocellulosic biomass-techniques, applications and commercialization. INTECH,

Chennai, pp 251–275

- Mrudula, S., & Shyam, N. (2012). Immobilization of *Bacillus megaterium* MTCC 2444 by caalginate entrapment method for enhanced alkaline protease production. *Brazilian Archives of Biology and Technology*, 55(1), 135–144. doi:10.1590/S1516-89132012000100017
- Nedovic, V., Gibson, B., Mantzouridou, T. F., & Bugarski, B. (2014). Aroma formation by immobilized yeast cells in fermentation processes. *Biochemistry*, 32(1), 173–216. doi: 10.1002/yea.3042
- Ni, Y., & Chen, R. (2009). Extracellular recombinant protein production from *Escherichia coli*. *Biotechnology Letters*, 31(11), 1661–1670. doi: 10.1007/s10529-009-0077-3
- Niu, X., Wang, Z., Li, Y., Zhao, Z., Liu, J., Jiang, L., Li, Z. (2013). “Fish-in-net”, a novel method for cell immobilization of *Zymomonas mobilis*. *PloS One*, 8(11), e79569. doi:10.1371/journal.pone.0079569
- Nkohla, A., Okaiyeto, K., Olaniran, A. & Nwodo, U. (2017). Optimization of growth parameters for cellulase and xylanase production by *Bacillus* species isolated from decaying biomass. *Journal of Biotech Research*, 8, 33–47.
- Nor, S., Azaman, A., Ramanan, R., & Tan, J. (2010). Screening for the optimal induction parameters for periplasmic producing interferon- α 2b in *Escherichia coli*. *African Journal of Biotechnology*, 9(38), 6345–6354.
- Notley, S. M., Russell J. C., and Elena P. I. (2016). Bacterial Interaction with Graphene Particles and Surfaces. *Science, Technology and Medicine*: 1–14. doi: 10.5772/56172
- Norsyahida, A., Rahmah, N., & Ahmad, R. M. Y. (2009). Effects of feeding and induction strategy on the production of BmR1 antigen in recombinant *E. Coli*. *Applied Microbiology*, 49, 544–50. doi: 10.1111/j.1472-765X.2009.02694.x
- Papaneophytou, C. P., & Kontopidis, G. (2014). Statistical approaches to maximize recombinant protein expression in *Escherichia coli*: A general review. *Protein Expression and Purification*, 94, 22–32. doi.org/10.1016/j.pep.2013.10.016.
- Patel, H., & Gupte, A., G. (2016). Optimization of different culture conditions for enhanced laccase production and its purification from *Tricholoma giganteum* AGHP. *Bioresourcing and Bioprocessing*. 3(11), 1–10. doi: 10.1186/s40643-016-0088-6
- Peniophora, S., Dinh, K. T. & Dinh, T. Q. (2013). Optimization of culture conditions and medium components for carboxymethyl cellulase (CMCase) production by a novel

- basidiomycete strain *Peniophora* sp. NDVN01. *Iranian Journal of Biotechnology*, 11(4): 251–259. doi: 10.5812 / ijb.11039
- Pervin, M. S., Jahan, M. G., Rana, A. Y. & Rahman, M. (2013). Effects of some environmental variables on urease in germinating chickpea (*Cicer arietinum* L.) seed. *Journal of Stress Physiology & Biochemistry*, 9(3), 345–356.
- Pithadiya, D., Deep, N., & Aarti, T. (2016). “Partial Purification and Optimization of Xylanase from *Bacillus Circulans*.” *Archives of Applied Science Research* 8(2): 1–10.
- Poet, P. D., Yves, A., & Roger, B. (1987). Plasmid stability in immobilized and free: recombinant *Escherichia coli* JM105 (pKK223-200): Importance of oxygen diffusion, growth rate, and plasmid copy number. *Applied and Environmental Microbiology*, 53(7), 1548–55.
- Radwan, H. H., Ihab, M. M., & Ibrahim, A. A. (2011). Optimization of a fed-batch fermentation process for production of bleomycin by *Streptomyces mobaraensis* ATCC 15003. *African Journal of Biotechnology*, 10(9), 1690–1695. doi: 10.5897/AJB10.1671
- Rahman, A. N. H. *Xylitol production of recombinant Escherichia coli immobilized on multi walled carbon nanotubes*. Master thesis, Universiti Teknologi Malaysia; 2016
- Ramanjaneyulu, H. N., Prasanna, G., & Rajasekhar, R. (2016). Optimization of cellulase production by *Penicillium* Sp. *Biotech*, 6(162): 1–11. doi: 10.1007/s13205-016-0483-x
- Ramchuran, S. O., Olle, H., & Nordberg, K. (2005). Effect of post induction nutrient feed composition and use of lactose as inducer during production of thermostable xylanase in *Escherichia coli* glucose-limited fed-batch cultivations. *Journal of Bioscience and Bioengineering*, 99(5): 477–84. doi: 10.1263/jbb.99.477
- Ramírez-Nuñez, J., Romero-Medrano, R., Nevárez-Moorillón, G. V., & Gutiérrez-Méndez, N. (2011). Effect of pH and salt gradient on the autolysis of *Lactococcus lactis* strains. *Brazilian Journal of Microbiology*, 52, 1495–1499.
- Robitaille, J., Chen, J., & Jolicoeur, M. (2015). A single dynamic metabolic model can describe mAb producing CHO cell batch and fed-batch cultures on different culture media. *Biotechnology Bioengineering*, 10(9), 1-23
- Rosano, G. L., & Ceccarelli, E. A. (2014). Recombinant protein expression in *Escherichia coli*: advances and challenges. *Frontiers in Microbiology*, 5(172), 1–17. <http://doi.org/10.3389/fmicb.2014.00172>

- Rostro, A., Magdalena, D. J., Mancera, A. E. I., & Beatriz, M. (2016). Nanobiocatalysis: nanostructured materials: a minireview. *Biocatalysis*, 2(1): 1–24. doi: 10.1515/boca-2016-0001
- Rusli, F. M., Mohamed, M. S., Mohamad, R., & Ni Nyoman Tri Puspaningsih, & Arbakariya B. Ariff. (2009). Kinetics of xylanase fermentation by recombinant *Escherichia coli* DH5 α in shake flask culture, *American Journal of Biochemistry and Biotechnology*, 5(3), 109–117.
- Sabu, A., Kiran, G. S., & Pandey, A. (2005). Purification and characterization of tannin acyl hydrolase from *Aspergillus niger* ATCC 16620, *Food Technology and Biotechnology*, 43(2), 133–138.
- Saifuddin N., Raziah A. Z., & Junizah A. R. (2016). Carbon nanotubes: A review on structure and their interaction with proteins, *Journal of Chemistry*, 2013(2013), 1–16.
- Salamanca C., L., Ashe, C. S., Stipanovic, A. J., & Nomura, C. T. (2014). Enhanced production of polyhydroxyalkanoates (PHAs) from beechwood xylan by recombinant *Escherichia coli*. *Applied Microbial and Cell Physiology*, 831–842.
- Salwanee, S., Aida, W. M. W., Mamot, S., Maskat, M. Y., & Ibrahim, S. (2013). Effects of enzyme concentration, temperature, pH and time on the degree of hydrolysis of protein extract from viscera of tuna (*Euthynnus affinis*) by using alcalase, *Sains Malaysiana*, 42(3), 279–287
- Schutter, K., Lin, Yao, C., Tiels, P., & Hecke, A. (2009). genome sequence of the recombinant protein production host *Pichia pastoris*. *Nature Biotechnology*, 27(6): 561–88. doi: 10.1038/nbt.1544
- Sezonov, G., & Ari, R. D. (2007). *Escherichia coli* physiology in Luria-Bertani Broth, *Bacteriology*, 189(23), 8746–8749.
- Shahi, N., Hasan, A., Akhtar, S., Siddiqui, M. H., Sayeed, U., Khan, M. K. A. (2016). “Xylanase : A Promising Enzyme.” *Journal of Chemical and Pharmaceutical Research* 8(3): 334–39.
- Shalinimol, C. R. (2016). A study on optimization of microbial alpha-amylase production. *International Journal of Advances in Microbiology & Microbiological Research*, 1(1), 22–27.
- Shan, G., Xing, J., Zhang, H., & Liu, H. (2015). Biodesulfurization of dibenzothiophene by microbial cells coated with magnetite nanoparticles, *Applied and Environmental Microbiology*, 71(8), 1–8. doi:10.1128/AEM.71.8.4497-4502.2005

- Shang, T., Si, D., Zhang, D. & Liu, X. (2017). Enhancement of thermoalkaliphilic xylanase production by *Pichia pastoris* through novel fed-batch strategy in high cell-density fermentation. *BMC Biotechnology*, 17(55), 1–10.
- Silva, L. C. A., Honorato, T. L., Cavalcante, R. S., Franco, T. T., & Rodrigues, S. (2016). Effect of pH and temperature on enzyme activity of chitosanase produced under solid stated fermentation by *Trichoderma* spp. *Indian Journal of Microbiology*, 52(1), 60–65. doi:10.1007/s12088
- Supelco, S. A. (1998). Guide to Solid Phase Extraction, Supelco Chemicals, Fluka Manual Procedure
- Suzana, C. S. M., Claudia, M., M., Larissa M. C. G. F., & Sandra, T. S. (2013). Immobilization of microbial cells: A promising tool for treatment of toxic pollutants in industrial wastewater. *African Journal of Biotechnology*, 12(28), 4412–4418. doi:10.5897/AJB12.2677
- Terrasan, C. R. F., José, M. G., & Eleonora, C. C. (2016). “Xylanase and β -Xylosidase from *Penicillium Janczewskii* : Puri Fi Cation, Characterization and Hydrolysis of Substrates.” *Electronic Journal of Biotechnology*: 1–10.
- Thirumarimurugan, M. (2016). Methods of cell immobilization and its application, *International Journal of Innovative Research in Science, Engineering and Technology*, 5(4), 5429–5433. doi:10.15680/IJIRSET.2016.0504175
- Tripathi, Nagesh, K. (2016). “Production and Purification of Recombinant Proteins from *Escherichia Coli*.” *ChemBioEng Rev* 3(3): 116–33. doi: 10.1002/cben.201600002
- Vélez, A. M., Carlos, A. & Horta, L. (2013). “Enhanced Production of Recombinant Thermo-Stable Lipase in *Escherichia Coli* at High Induction Temperature.” *Protein expression and purification* 90: 96–103.
- Vengadaramana, A., Balakumar, S., & Vasanthy, A. (2014). Effect of temperature, pH, substrate (starch) and glucose on stability of α - amylase from *Bacillus licheniformis* ATCC 6346. *Scholars Academic Journal of Pharmacy*, 3(6), 492–95.
- Waluyo, J., Dian, B., Nurul, H., & Yanni, S. (2017). “Immobilization of *Saccharomyces Cerevisiae* Using Ca- Alginate for Bioethanol Production from Empty Fruit Bunch of Oil Palm.” In American Institute of Physics Conference Proceedings, 020016–1–020016–18.
- Walia, A., Shiwani, G., Preeti, M., & Anjali, C. (2017). “Microbial Xylanases and Their Industrial Application in Pulp and Paper Biobleaching : A Review.” *Biotech* 7(11): 1–12. doi: 10.1007/s13205-016-0584-6

- Wang, J., Yangyuan, L., & Danni, L. (2016). Improved production of *Aspergillus usamii* endo- β -1,4-xylanase in *Pichia pastoris* via combined strategies. *Biomed Research International*, 1–9. doi: org/10.1155/2016/3265895
- Ward, O. P. (2012). “Production of Recombinant Proteins by Filamentous Fungi.” *Biotechnology Advances* 30: 1119–39. doi: 10.1016/j.biotechadv.2011.09.012
- Werbowy, O., Sławomir, W., & Tadeusz, K. (2017). Plasmid stability analysis based on a new theoretical model employing stochastic simulations. *Plos One* 12(8), 1–21. doi: 10.1371/journal.pone.0183512
- Wyre, C., & Overton., T. W. (2014). Use of a stress - Minimisation paradigm in high cell density fed - batch *Escherichia coli* fermentations to optimise recombinant protein production. *Journal of Microbiology Biotechnology*, 41, 1391–1404. doi: 10.1007/s10295-014-1489-1
- Xu, J., Sun, J., Wang, Y., Sheng, J., Wang, F., & Sun, M. (2014). Application of iron magnetic nanoparticles in protein immobilization, *Molecules*, 19(8), 11465–11486. doi:10.3390/molecules190811465.
- Yates, G. T., & Smotzer, T. (2007). The lag phase and initial decline of microbial growth curves. *Journal of Theoretical Biology*. 244, 511 – 517.
- Yavari, R. (2015). Adsorption of cesium (I) from aqueous solution using oxidized multiwall carbon nanotubes, *Journal of Radioanalytical and Nuclear Chemistry*, 287(2), 1–22.
- Yoon, S. H., Kim, S. K., & Kim, J. F. (2010). Secretory production of recombinant proteins in *Escherichia coli*, *Recent Patents on Biotechnology*, 4(1), 23–29.
- Zajkoska, P., Martin R., & Michal R. (2013). Biocatalysis with immobilized *Escherichia coli*. *Applied Microbiology and Biotechnology*, 97, 1441–1455. doi: 10.1007/s00253-012-4651-6
- Zhang, D., Yuwen, L., & Peng, L. (2016). Parameters affecting the performance of immobilized enzyme, *Journal of Chemistry*, 2013, 1–7.
- Zhang, J., Zhang, J., Zhang, F., Yang, H., Huang, X., Liu, H., & Guo, S. (2010). Graphene oxide as a matrix for enzyme immobilization. *Langmuir*, 26(9), 6083–6085. doi:10.1021/la904014z
- Zhang, L., Liang, J., Huang, Y. & Ma, Y. (2009). Size-controlled synthesis of graphene oxide sheets on a large scale using chemical exfoliation. *Nanoscale Research Letters*, 7(47), 3365 –3380. doi: 10.1016/j.carbon.2009.07.045

- Zhang, P., Zhou, W., Wang, P., Wang, L., & Tang, M. (2013). Enhancement of chitosanase production by cell immobilization of *Gongronella* sp. JG. *Brazilian Journal of Microbiology*, 44(1), 189-195.
- Zhao, L., Geng, J., Guo, Y. & Liao, X. (2015). Expression of the thermobifida fusca xylanase xyn11a in *pichia pastoris* and its characterization. *BMC Biotechnology*, 15(18), 1–12. doi: 10.1186/s12896-015-0135-y

